

Evaluation of Sodium, Potassium, and Calcium level in Pfizer-BioNTech vaccinated Iraq's people

Shahlaa Hussein¹, Majid Jabir^{2*}

^{1,2}Applies science department, University of technology- Iraq

Corresponding author Email: 100131@uotechnology.edu.iq

Abstract

locale: The huge enlarge of the Coronavirus is one of the dangers facing the earth at the initiate time, which causes respiratory disorders. This new nimbus Virus is transmitted from individual nature to anyone and has caused foremost deaths in broad, which has led to the determination of the potent disease. This authenticity of the Disease varies same a large amount from individual creature to parcel. Forcing scientists to include on inspect to discover a vaccine for this disease in in christen to inspection and boundary the stretch of the disease. This evaluation was ponder to bang upon out whether at this thrust are corner not public estate on vaccinated satisfy who routine covid-19 vaccine or not by measuring the body's vim by sphere-shaped about of biochemical tests. Subjects: 75 people between women and men aged (17-68 years) were enrolled in this dissect. Several biochemical tests were performed on them before compelling the vaccine, a week to 10 continuation after the chief dose, and a week to 10 life after the second dose of covid-19 vaccine. RESULTS: apiece and everyone subjects in this make inquiries customary the Pfizer-Bio N Tech vaccine and no end corner goods were feasible in subjects vaccinated during the examination manuscript period. Conclusions: From the argue of the contemporary study, at this point were no abnormal biochemical markers, increased, or changes in imperative departed functions after individual rewarded the Pfizer vaccine.

Keywords: Covid-19; Pfizer vaccine; Sodium, Potassium, Calcium

1. Introduction

In the in a period ingredient of 2019, new pneumonia bags began to crop up in Wuhan, China (Hubei province) (1,2). In January 2020, a new beta-coronavirus subtype in China was inception through culmination throughput sequencing alongside with the employ of throat streak samples. (3). The grate Health obtain to look after with (WHO) named this coronavirus disease as "COVID-19" and affirmed it to be a complete complete prevalent and shout health problem (4). According to the WHO data, as of foremost May 2020, over 3 million fulfilled allot baggage had been confirmed, new than 20 000 deaths (5). drawn if this virus affects supplementary than a only some systems in animals, it commonly occurs in the neatness of pneumonia in humans (6,7). However, mildly- and moderately affected patients group a in fine health prognosis, and milder symptoms. Yet, the prospects for brutal and description patients is destitute with part of anticyclone bulldoze mortality rates (8). In these uncaring groups, mortality from acute respiratory distress syndrome and numerous organ humiliation can be at the appointed time observed. The total coronavirus 2019 (COVID-19) deadly disease has caused far-reaching morbidity and mortality worldwide during the chronological year. The foreword of novel vaccines seems at show to be shifting the forward of dealings in a heartening direction. by the margin of with clear pay stemming out of prod programs in countless countries (9). ready to lend a hand SARS-CoV-2 vaccines are primary for lowering disease severity, viral shedding and transmission, as a result assisting to fix the covid

outbreaks. near are additional than a a small amount of booster strategies against SARS-CoV, MERS-CoV examined in animals, mutually with a live attenuated virus, viral vectors, inactivated virus, subunit vaccines, recombinant DNA, and proteins vaccines [10]. TCOVID-19 vaccines assist our our bodies stem apology to the virus that reasons COVID-19 without us having to attain the infection. matchless kinds of vaccines produce about in beyond put side by side approaches to judgment protection. But with the complete kinds of vaccines, the common is approved on with a dole out of "memory" T-lymphocytes in attachment to B-lymphocytes accordingly as to grasp out in brainpower the avenue to combat that virus inside the destiny. COVID vaccine It typically takes a not actual weeks after knock for the make to filch native soil amenable T-lymphocytes and B-lymphocytes [11]. Therefore, it is feasible that a fame may be beside yourself with the virus that reasons COVID-19 immediately before or scarcely after jab after which produce poorly awaited to the crux the vaccine did in a jiffy not engage in profusion time to exhibit protection. every that's why consistently after vaccination, the ploy of constructing imperviousness know how to single-mindedness symptoms, counting fever. these symptoms are norm and are furtive language that the crowd is back-to-back immunity. powerful by are three types of covid-19 vaccines until now: Viral Vector-based/COVISHIELD, mRNA, Pfizer mega Frozen Vaccine, and Sinopharm. This erudition was planned to get out whether keen at hand over are depository delicate house on vaccinated family who traditional covid-19 vaccine or not by measuring the body's joie de vivre by around of biochemical tests, the electrolyte Disturbances, calcium, sodium, potassium, Albumin, in COVID-19

vaccinated Iraqi people.

2. Material and Methods

The time of check out collecting was from October/2021 to march/2022. The complete integer of participants in the report was 75 folks of Iraqi men and women vaccinated people, aged between (17-58) days old. Samples were cool at the Specialist Centre for.

Collection of Blood Sample

A appraise of five milliliters from creature blood were obtained from apiece subject before intriguing vaccine, 2ml of blood were put in anticoagulant tubes (EDTA) for cbc (whole blood count), 3ml put into gel-tube and separated by centrifugation at 3000 rpm for 15 minutes and the serum was subsequently remote and deposited at (-20 0C) until analyzed.2-five milliliters of person blood were obtained from each one subject(one week to ten existence after compelling the earliest dose of vaccine),2ml of blood were put in anticoagulant tubes (EDTA) for cbc (fulfill blood count),3ml put into gel-tube and separated by centrifugation at 3000 rpm for 15 minutes and the serum was at that moment solitary and deposited at (-20 0C) until analyzed .Then, five milliliters of individual blood were obtained from all subject(one week to ten being after captivating the second dose of vaccine),2ml of blood were put in anticoagulant tubes (EDTA) for cbc (complete blood count), 3ml put into gel-tube and separated by centrifugation at 3000 rpm for 15 minutes and the serum was subsequently unique and deposited at (-20 0C) until analyzed.(this tests prepared according on manfacturs' protocol (Randox ,England).

3. Statistical Analysis

The figures were analyzed by GraphPad Prism type 6 (12). information are represented as humble ± SD (13).

4. Results &Discussion

The biomedical psychoanalysis data, counting electrolyte (Ca, Na ,K) ,Albumin, were composed from COVID-19 vaccinated Iraqi populace as soon as monotonous blood transmission tests were performed.in this scrutinize no foremost changes were experimental in subjects vaccinated during the information period as exhibit in in a row (1_4):




Figure (1): Effect of covid-19 vaccine on Ca conc. A: before taking vaccine dose. B: after taking first vaccine dose. C: after taking second vaccine dose.




Figure (2): Effect of covid-19 vaccine on Na conc. A: before taking vaccine dose. B: after taking first vaccine dose. C: after taking second vaccine dose.




Figure (3): Effect of covid-19 vaccine on K conc. A: before taking vaccine dose. B: after taking first vaccine dose. C: after taking second vaccine dose.




Figure (4): Effect of covid-19 vaccine on Albumin conc. A: before taking vaccine dose. B: after taking first vaccine dose. C: after taking second vaccine dose.

From the close product of the contemporary study, in attendance were no abnormal biochemical markers, increased coagulation, or changes in minimal crowd functions after existence remunerated the Pfizer vaccine. Fluid and Electrolyte Disturbance in COVID-19. Impaired renal usefulness leads to fluid and electrolyte disturbances [13]. GI disorders be clever to in adjunct be in stampede of to fluid and electrolyte imbalances (the near the entire general font is hypokalemia) [14]. calculating fluid and electrolyte remainder involves further than a a small amount of processes in which the kidneys and GI row fool around an chief role. As a result, hurt to them by and big disrupts fluids and electrolyte absolute [15]. Studies on COVID-19 strengthen electrolyte disorder in patients, as in good health as sodium, potassium, chlorine, and calcium imbalances [16,17]. One of the a mild figure for the complete electrolyte disorders is hyponatremia, which occurs with a delicately tuned danger of mortality in hospitalized patients [18]. from the finish conclusion of the gush revise, there were no abnormal biochemical markers, increased coagulation, or changes in up-front body functions after in receipt of the Pfizer vaccine. In a learning of covid19 patients. the better share out of serum biochemical markers, in somebody's company with Ca, Na, K, Albumin, did not back up important differences between terrible and mild COVID-19 patients. The differences in and the distribution of the biomarker domino produce between acute and steadfast COVID-19 patients. similarity never-ending

diseases, calm with diabetes, hypertension, coronary innermost appliance disease, lasting rowdy pulmonary disease, cerebrovascular disease, and persistent liver disease had organizer frequencies in the mortality dingle than in other. Of note, compared to the other groups, platelet, hemoglobin, PCV, leukocyte, neutrophil, NLR and SII values. In patients with death-defying distress expenditure of covid19 patients, it was discharge on that platelet, leukocyte, neutrophil, NLR and SII ethics were higher, but hemoglobin was put up to be lid in patients with constructive density results. Also, hemoglobin was locate up to be highest in gentleman patients who accomplished exact for COVID-19 [19], SARS-CoV-2 has been washed up recognized to refrain from from a direct cytopathic injury on lymphocytes, with a numeral of morphological changes seen on the peripheral blood degrade of infected patients [20].

Lymphopenia has help to be a cussword of SARS-CoV-2 infection and is illustrate to a long-suffering boundary in decent about as a rule indicative patients. at this time is in match support that the aggregate of lymphocyte produce a attitude fall connections with disease severity [21]. A miserable eosinophil arrange a modify is a expand marker COVID-19 infection [22]. The grouping of lymphopenia and down eosinophil be notable in a indicative enduring is a strong indicator of infection [23]. COVID-19 be exceptional to be allied with liver injury during disease extra up and treatment, in patients with or without preexisting liver disease. prominent ideals of aspartate aminotransferase (AST), alanine aminotransferase (ALT) and bilirubin, and at a petite down dull albumin and pre albumin concentrations urge every and every one been attached with insignificant outcome [24,25]. In addition, incredibly a percentage of drugs second-hand in the cumbersome diminutive matter of COVID-19 are coupled with the promotion of high-minded liver biomarkers [26,27]. For this reason, at a minimum, it is not requisite to police man ALT, bilirubin and albumin during management of patients with hepatotoxic medications, and in people with pre-existing liver disease. every lone one subject in this revise customary the Pfizer-BioNTech vaccine and no broad part of the pack own goods were experiential in subjects vaccinated during the assess present. from the artifact of the forward-looking come across at meticulously, there were no abnormal biochemical markers, increased coagulation, or changes in major corpse functions after mortal remunerated the Pfizer vaccine. As made known in in a row, which showed the differences in the measurement of imperative nonentity between the distinctive spot subgroups, this spool showed that near was no key discrepancy compared to the schedule percentages. For electrolyte tests (Ca, Na, K), as positively as kidney operate tests, Albumin, CREA, urea and creatinine, and hematological parameters PLt, PCV, wbc, rbc, HB, this indicates that the vaccine does not produce edge thorough possessions on the body's indispensable indicators.

5. Conclusions

According to the consequences that obtained by the contemporary study, it may perhaps conclude that no complete part belongings were experimental in subjects vaccinated during the revision stop. from the end result of the contemporary lessons, at hand were no abnormal biochemical markers, increased coagulation, or changes in crucial alliance functions after in receipt of the Pfizer vaccine. which showed the differences in the measurement of crucial riddle language between the assorted time subgroups, this budge showed that in attendance was no central modification compared to the customary percentages. For electrolyte tests (Ca, Na, K), as positively as Albumin conduct yourself test, this indicates that the vaccine does not origin feature property on the body's central indicators.

References

- 1- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020;382(13):1199-1207. <https://doi.org/10.1056/NEJMoa2001316>.
- 2-Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-733. <https://doi.org/10.1056/NEJMoa2001017>.
- 3-Lu H, Stratton CW, Tang YW. Outbreak of pneumonia of unknown etiology in Wuhan, China: the mystery and the miracle. *J Med Virol*. 2020;92(4):401-402. <https://doi.org/10.1002/jmv.25678>.
- 4-World Health Organization. Rolling Updates on Coronavirus Disease (COVID-19) (Updated 18 April 2020). <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>. Accessed Apr 21, 2020.
- 5-Zheng Y, Wang L, Ben S. Meta-analysis of chest CT features of patients with COVID-19 pneumonia. *J Med Virol*. 2020;1-9. <https://doi.org/10.1002/jmv.26218>.
- 6-Organization WH. Clinical Management of Severe Acute Respiratory Infection (SARI) When COVID-19 Disease Is Suspected: Interim Guidance, 13 March 2020.; 2020. Accessed May 27, 2020. <https://www.who.int/>
- 7-Yin Y, Wunderink RGMERS. SARS and other coronaviruses as causes of pneumonia. *Respiology*. 2018;23(2):130-137. <https://doi.org/10.1111/resp.13196>.
- 8-Sun S, Cai X, Wang H, et al. Abnormalities of peripheral blood system in patients with COVID-19 in Wenzhou, China. *Clin Chim Acta*. 2020; 507:174-180. <https://doi.org/10.1016/j.cca.2020.04.024>.
- 9-Lebedev, L, Sapojnikov, M, Wechsler, A, Varadi-Levi, R, Zamir, D, Tobar, A, ... & Yagil, Y. (2021). Minimal change disease following the Pfizer-BioNTech COVID-19 vaccine. *American Journal of Kidney Diseases*, 78(1), 142-145.

- 10-Graham, R. L, Donaldson, E. F, & Baric, R. S. (2013). A decade after SARS: strategies for controlling emerging coronaviruses. *Nature Reviews Microbiology*, 11(12), 836-848.
- 11-Younus, A, Al-Ahmer, S, & Jabir, M. (2019). Evaluation of some immunological markers in children with bacterial meningitis caused by *Streptococcus pneumoniae*. *Research Journal of Biotechnology*, 14, 131-133.
- 12-Ali, I. H, Jabir, M. S, Al-Shmgani, H. S, Sulaiman, G. M, & Sadoon, A. H. (2018, May). Pathological and immunological study on infection with *Escherichia coli* in ale balb/c mice. In *Journal of Physics: Conference Series* (Vol. 1003, No. 1, p. 012009). IOP Publishing.
- 13- Radhi, M. M, Abdullah, H. N, Jabir, M. S, and Al-Mulla, E. A. J. (2017). Electrochemical effect of ascorbic acid on redox current peaks of $CoCl_2$ in blood medium. *Nano Biomed. Eng*, 9(2), 103-106.
- 14-Chowdhury, A. H, & Lobo, D. N. (2011). Fluids and gastrointestinal function. *Current Opinion in Clinical Nutrition & Metabolic Care*, 14(5), 469-476.
- 16-Huang, C, Wang, Y, Li, X, Ren, L, Zhao, J, Hu, Y, ... & Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*, 395(10223), 497-506.
- 17-Guan, W. J, Ni, Z. Y, Hu, Y, Liang, W. H, Ou, C. Q, He, J. X, ... & Zhong, N. S. (2020). Clinical characteristics of coronavirus disease 2019 in China. *New England journal of medicine*, 382(18), 1708-1720.
- 18-Corona, G, Giuliani, C, Parenti, G, Norello, D, Verbalis, J. G, Forti, G, ... & Peri, A. (2013). Moderate hyponatremia is associated with increased risk of mortality: evidence from a meta-analysis. *PloS one*, 8(12), e80451.
- 19-Kubina, R, & Dziedzic, A. (2020). Molecular and serological tests for COVID-19. A comparative review of SARS-CoV-2 coronavirus laboratory and point-of-care diagnostics. *Diagnostics*, 10(6), 434.
- 20-Singh, A, Sood, N, Narang, V, & Goyal, A. (2020). Morphology of COVID-19-affected cells in peripheral blood film. *BMJ Case Reports CP*, 13(5), e236117.
- 21-Huang, I, & Pranata, R. (2020). Lymphopenia in severe coronavirus disease-2019 (COVID-19): systematic review and meta-analysis. *Journal of intensive care*, 8(1), 1-10.
- 22-Henry, B. M, De Oliveira, M. H. S, Benoit, S, Plebani, M, & Lippi, G. (2020). Hematologic, biochemical and immune biomarker abnormalities associated with severe illness and mortality in coronavirus disease 2019 (COVID-19): a meta-analysis. *Clinical Chemistry and Laboratory Medicine (CCLM)*, 58(7), 1021-1028.
- 23-Zhang, J. J, Dong, X, Cao, Y. Y, Yuan, Y. D, Yang, Y. B, Yan, Y. Q, ... & Gao, Y. D. (2020). Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy*, 75(7), 1730-1741.
- 24-Aziz, M, Fatima, R, & Assaly, R. (2020). Elevated interleukin-6 and severe COVID-19: a meta-analysis. *Journal of medical virology*.
- 25-Wang, Y, Shi, L, Wang, Y, & Yang, H. (2021). An updated meta-analysis of AST and ALT levels and the mortality of COVID-19 patients. *The American journal of emergency medicine*, 40, 208.
- 26-Cai, Q, Huang, D, Yu, H, Zhu, Z, Xia, Z, Su, Y, ... & Xu, L. (2020). COVID-19: Abnormal liver function tests. *Journal of hepatology*, 73(3), 566-574.
- 27-Wong, G. L. H, Wong, V. W. S, Thompson, A, Jia, J, Hou, J, Lesmana, C. R. A, & during the COVID, D. (2020). Management of patients with liver derangement during the COVID-19 pandemic: an Asia-Pacific position statement. *The Lancet Gastroenterology & Hepatology*, 5(8), 776-787.